arXiv:math/0610823v1 [math.QA] 27 Oct 2006

A GENERALIZATION OF COXETER GROUPS, ROOT SYSTEMS, AND MATSUMOTO'S THEOREM

I. HECKENBERGER AND H. YAMANE

Abstract. The root systems appearing in the theory of Lie superalgebras and Nichols algebras admit a large symmetry extending properly the one coming from the Weyl group. Based on this observation we set up a general framework in which the symmetry object is a groupoid. We prove that in our context the groupoid is generated by reflections and Coxeter relations. This answers a question of Serganova. Our weak version of the exchange condition allows us to prove Matsumoto's theorem. Therefore the word problem is solved for the groupoid.

1. Introduction

Kac-Moody algebras [Kac90] and their generalizations enjoy continuously growing interest since their introduction in the late sixties. Lie superalgebras [Kac77] became central objects in physical models, and generalized Kac-Moody-Borcherds algebras [Bor88] turned out to be important in group theory in connection with the monster. One of the basic features of these algebras is their relation to root systems, Weyl groups, and their generalizations.

In connection with contragredient Lie superalgebras Serganova [Ser96] found out that the Weyl group symmetry of the Lie algebra can be extended using reflections on isotropic roots. These so called "odd reflections" were used by the second author [Yam99] to analyze the structure of affine super Lie algebras very efficiently. Serganova [Ser96] gave an axiomatic definition of "generalized root systems" (GRS) which is based on ordinary and odd reflections, and classified them completely. Below Example 6.7 she writes the following.

- "It is an interesting question what is the analogue of Weyl group for GRS. It is natural to consider the finite group $\widetilde{W}$ generated by s_α for all $\alpha \in R$. This group in general is not a subgroup of $\mathrm{GL}(V)$. It is unknown if this group is Coxeter."

Our paper gives an answer to this question as explained in Remark 25.

The discovery of quantized Kac-Moody algebras by Drinfel'd [Dri87] and Jimbo [Jim86] opened a new major research direction with many novel features, ideas, and connections to other research fields. Motivated by these examples, Andruskiewitsch and Schneider [AS98] launched a project to classify pointed Hopf algebras with certain finiteness conditions, and performed it successfully under some hypotheses [AS05]. Their method is based on the knowledge of the finiteness property of Nichols algebras, the latter being main ingredients for the construction of quantized Kac-Moody algebras. These finiteness properties can be determined effectively using the Weyl groupoid [Hec06b] attached to a Nichols algebra. Also the wish to understand better the structure of this groupoid gave motivation for our work.

In the literature there exist several generalizations of root systems, mostly depending on the applications. A very nice general treatment with many references is the work of Loos and Neher [LN05]. In our paper we do not attempt to give a complete set of axioms for a root system. Instead we concentrate on its property being invariant under symmetry. Motivated by Serganovas work and the appearence of the Weyl groupoid for Nichols algebras we require a large symmetry in the sense that reflections with respect to all simple roots should be defined. Unfortunately this excludes already from the beginning the root systems of generalized Kac-Moody-Borcherds algebras. However Kac-Moody algebras and a very large class of their super and quantized analogues are still covered, together with so far less understood examples appearing in the classification of Nichols algebras of diagonal type [Hec06a]. Our main results are Theorems 24, 27 and 29. Theorem 24 gives an answer to Serganovas question about the right generalization of the Weyl group (see also Remark 25). The groupoid attached naturally to the root system of a Lie superalgebra is a Coxeter groupoid. The result holds more generally for any root system satisfying the axioms required in Definition 5. Further, Coxeter groups are known to satisfy the exchange condition. In our generality this property does not hold in its standard form. However Theorem 27 gives a weak version of it, which seem to be sufficient in many applications. In particular we are able to prove Matsumoto's theorem [Mat64], see Theorem 29, which states that any two reduced expressions of the same length can be transformed to each other using Coxeter relations only. This means that if the groupoid is finitely generated then with finite calculation it is possible to check whether two given elements in the groupoid are equal. Hence the word problem is solved for any groupoid appearing in Theorem 29. Finally, in the last section of the paper we give an example which appears in the classification of Nichols algebras but is related neither to a Kac-Moody nor to a Lie superalgebra.

In this paper let $\mathbb{N}$ denote the set of positive integers, and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. The composition sign for general groups will usually be omitted.

2. Group actions

For any set N let $F(N)$ denote the free group generated by the elements of N and their inverses. Further, let $F_2(N)$ denote the free group generated by the elements of N as involutions. With other words, $F_2(N)$ is the quotient of $F(N)$ by the subgroup consisting of products of elements gn^2g^{-1} and $gn^{-2}g^{-1}$, where $g \in F(N)$ and $n \in N$. Since in $F_2(N)$ one has $n = n^{-1}$ for all $n \in N$, $F_2(N)$ is generated as a monoid by the elements of N. The unit of $F_2(N)$ is denoted by 1.

Let N and A be nonempty sets. An action $\triangleright$ of $F_2(N)$ on A is a map $\triangleright : F_2(N) \times A \to A$ such that $1 \triangleright a = a$ and $g \triangleright (h \triangleright a) = (gh) \triangleright a$ for all $g, h \in F_2(N)$ and $a \in A$. By definition of $F_2(N)$ this is equivalent to

$$n \triangleright (n \triangleright a) = a \qquad \text{for all } n \in N, a \in A. \tag{1}$$

An action $\triangleright : F_2(N) \times A \to A$ is called *transitive*, if for each $a, b \in A$ there exists $g \in F_2(N)$ such that $g \triangleright a = b$. For an arbitrary action $\triangleright : F_2(N) \times A \to A$ one has a unique decomposition of A into a disjoint union of subsets A_p, such that $F_2(N) \triangleright A_p \subset A_p$ and the restriction of $\triangleright$ to $F_2(N) \times A_p$ is transitive for all p.

Let N be a nonempty set and $N' \subset N$ a nonempty subset. Then $F_2(N')$ can be regarded as a subgroup of $F_2(N)$. Let $\triangleright$ be an action of $F_2(N)$ on A. Then

$\triangleright$ induces an action $\triangleright' : F_2(N') \times A \to A$ which is called the restriction of $\triangleright$ to $F_2(N') \times A$. The transitivity of $\triangleright$ does not imply transitivity of $\triangleright'$.

Let $\triangleright : F_2(N) \times A \to A$ be an action and $n, n' \in N$. For each $a \in A$ define

$$\Theta(n, n'; a) := \{(nn')^m \triangleright a,\ (n'n)^m \triangleright a \,|\, m \in \mathbb{N}_0\}.$$

Let $\theta(n, n'; a) := |\Theta(n, n'; a)|$, the cardinality of $\Theta(n, n'; a)$, which is either in $\mathbb{N}$ or is ∞. One obviously has $\Theta(n, n'; a) = \Theta(n', n; a)$ and $\theta(n, n'; a) = \theta(n', n; a)$. Moreover, if $\theta(n, n'; a) \in \mathbb{N}$ then $\Theta(n, n'; a) = \{(nn')^m \triangleright a \,|\, 0 \le m < \theta(n, n'; a)\}$. Further, $\Theta(n, n'; n \triangleright a) = \Theta(n, n'; n' \triangleright a)$ by definition, and equations $\theta(n, n'; a) = \theta(n, n'; n \triangleright a) = \theta(n, n'; n' \triangleright a)$ hold.

Let $a_0 := a$, $b_0 := a$, and define recursively $a_{m+1} := n \triangleright b_m$, $b_{m+1} := n' \triangleright a_m$ for all $m \in \mathbb{N}_0$. Using Equation (1) one obtains that

$$\theta(n, n'; a) := \begin{cases} \infty & \text{if } a_m \ne b_m \text{ for all } m \in \mathbb{N}, \\ \min\{m \in \mathbb{N} \,|\, a_m = b_m\} & \text{otherwise.} \end{cases} \tag{2}$$

3. Coxeter groupoids

First we give a generalization of the notion of Coxeter groups to groupoids. For the notion of groupoids consult for example [CP61, Sect. 3.3].

Definition 1. Let N and A be nonempty sets and let $\triangleright$ be a transitive action of $F_2(N)$ on A. For each $a \in A$ and $i, j \in N$ with $i \ne j$ let $m_{i,j;a}$ be a multiple of $\theta(i, j; a)$ lying in $\mathbb{N} \setminus \{1\} \cup \infty$. Set $\mathbf{m} := (m_{i,j;a} \,|\, i, j \in N, i \ne j, a \in A)$. Let W be the groupoid generated by elements e_a and $s_{i,a}$, where $a \in A$ and $i \in N$, and the following relations:

$$\begin{aligned} e_a^2 &= e_a, & e_a e_b &= 0 \text{ for } a \ne b, \\ e_{i \triangleright a} s_{i,a} &= s_{i,a} e_a = s_{i,a}, & s_{i, i \triangleright a} s_{i,a} &= e_a, \end{aligned} \tag{3}$$

$$\begin{aligned} s_i s_j \cdots s_j s_{i,a} &= s_j s_i \cdots s_i s_{j,a} \ (m_{i,j;a} \text{ factors}) && \text{if } m_{i,j;a} \text{ is finite and odd,} \\ s_j s_i \cdots s_j s_{i,a} &= s_i s_j \cdots s_i s_{j,a} \ (m_{i,j;a} \text{ factors}) && \text{if } m_{i,j;a} \text{ is finite and even,} \end{aligned} \tag{4}$$

where a is running over all elements of A. The quintuple $(W, N, A, \triangleright, \mathbf{m})$ will be called a *Coxeter groupoid*.

Note that in our convention the multiplication of two elements in a groupoid is always defined, but the product may be the unique element 0 which satisfies the property $g0 = 0g = 0$ for all g. Moreover, in (4) and later on we use the following convention.

Convention 2. Let $(W, N, A, \triangleright, \mathbf{m})$ be a Coxeter groupoid, $(i_1, \ldots, i_m)$ a sequence of elements in N, and $(b_1, \ldots, b_m)$ a sequence of elements in A. By Relations (3) the element $w = s_{i_1,b_1} \cdots s_{i_{m-1},b_{m-1}} s_{i_m,b_m} \in W$ is obviously zero if one of the relations $b_j = i_{j+1} \triangleright b_{j+1}$, where $j \ge 1$, fails. Thus $w \ne 0$ implies that $b_j = i_{j+1} \cdots i_m \triangleright b_m$ for all $j < m$, and in the latter case we will write $s_{i_1} \cdots s_{i_{m-1}} s_{i_m,b_m}$ for w.

Define $\ell : W \to \mathbb{N}_0 \cup \{-\infty\}$ to be the function such that $\ell(0) = -\infty$, $\ell(e_a) = 0$ for all $a \in A$, and

$$\ell(w) = \min\{m \in \mathbb{N} \,|\, w = s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a} \text{ for some } i_1, \ldots, i_m \in N,\ a \in A\}$$

otherwise. One has

$$\ell(w) = \ell(w^{-1}), \tag{5}$$

$$\ell(ww') \leq \ell(w) + \ell(w') \tag{6}$$

for each $w, w' \in W$. One says that a product $w = s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a} \in W$ is *reduced*, if $m = \ell(w)$.

Note that if the cardinality of A is 1, then the action $\triangleright$ is necessarily trivial, and hence $\theta(i,j;a) = 1$ for $a \in A$ and all $i, j \in N$. Thus in this case the definition of $(W, N, A, \triangleright, \mathbf{m})$ coincides with the definition of a Coxeter group.

Remark 3. Coxeter groups are dealt with effectively using reflections obtained by conjugation of generators. Note that the structure constants $m_{i,j;a}$ of a Coxeter groupoid depend also on the elements of A, and hence in general the adjoint action of W on itself is not defined. Therefore the standard proofs can not always be generalized to our setting.

Remark 4. Coxeter groups satisfy the exchange condition, which follows from the existence of sufficiently many reflections. The exchange condition in its standard form does not hold for arbitrary Coxeter groupoids, see Section 7 for an example. A weak version of the exchange condition is given in Theorem 27 for a special class of Coxeter groupoids.

We end this section with the definition of special elements of W and giving some commutation rules which will be needed later.

Following the notation in Definition 1 assume that $a \in A$ and $i, j \in N$, where $i \neq j$. Define

$$C_{i,j;a} = \begin{cases} s_i s_j s_i \cdots s_i s_{j,a} \ (m_{i,j;a} - 1 \text{ factors}) & \text{if } m_{i,j;a} \text{ is odd,} \\ s_i s_j s_i \cdots s_j s_{i,a} \ (m_{i,j;a} - 1 \text{ factors}) & \text{if } m_{i,j;a} \text{ is even.} \end{cases} \tag{7}$$

Then Equation (4) implies the following relations.

$$s_i C_{j,i;a} = \begin{cases} C_{j,i:j \triangleright a} s_{j,a} & \text{if } m_{j,i;a} \text{ is odd,} \\ C_{j,i:i \triangleright a} s_{i,a} & \text{if } m_{j,i;a} \text{ is even.} \end{cases} \tag{8}$$

4. A GENERALIZATION OF ROOT SYSTEMS

This section is devoted to the definition of a generalization of root systems which admits the symmetry of a Coxeter groupoid. The required large symmetry parallels the main idea behind root systems of semisimple Lie algebras. In the examples given below we will show how classical objects fit into our definition.

Definition 5. Let $\mathcal{R}$ be the set consisting of all triples $(R, N, A, \triangleright)$ such that the following conditions hold.

1. N and A are sets and $\triangleright$ is a transitive action of $F_2(N)$ on A.
2. Let $V_0 = \mathbb{R}^{|N|}$. Then $R = \{(R_a, \pi_a, S_a) \,|\, a \in A\}$, where $\pi_a = \{\alpha_{n,a} \,|\, n \in N\} \subset R_a \subset V_0$, and π_a is a basis of V_0 for all $a \in A$.
3. $R_a = R_a^+ \cup -R_a^+$ for all $a \in A$, where $R_a^+ = \mathbb{N}_0 \pi_a \cap R_a$.
4. For any $i \in N$ and $a \in A$ one has $\mathbb{R}\alpha_{i,a} \cap R_a = \{\alpha_{i,a}, -\alpha_{i,a}\}$.
5. $S_a = \{\sigma_{i,a} \,|\, i \in N\}$, and for each $a \in A$ and $i \in N$ one has $\sigma_{i,a} \in \mathrm{GL}(V_0)$, $\sigma_{i,a}(R_a) = R_{i \triangleright a}$, $\sigma_{i,a}(\alpha_{i,a}) = -\alpha_{i,i \triangleright a}$, and $\sigma_{i,a}(\alpha_{j,a}) \in \alpha_{j,i \triangleright a} + \mathbb{N}_0 \alpha_{i,i \triangleright a}$ for all $j \in N \setminus \{i\}$.

(6) $\sigma_{i,i\triangleright a}\sigma_{i,a}=\mathrm{id}$ for $a\in A$ and $i\in N$.

(7) Let $a\in A$, $i,j\in N$, $i\neq j$, and $d=|\mathbb{N}_0\{\alpha_{i,a},\alpha_{j,a}\}\cap R_a|$. If d is finite then $\theta(i,j;a)$ is finite and it divides d.

The cardinality of N is called the *rank of* $(R,N,A,\triangleright)\in\mathcal{R}$.

For any $(R,N,A,\triangleright)\in\mathcal{R}$ and any $a\in A$ define

$$\begin{aligned}R^{\mathrm{re}}_a=\{\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,b}(\alpha_{j,b})\,|\,m\in\mathbb{N}_0,b\in A,\\ i_1,\dots,i_m,j\in N,i_1\cdots i_m\triangleright b=a\}.\end{aligned}\tag{9}$$

Similarly to Definition 5(3) we will use the notation $(R^{\mathrm{re}}_a)^+=\mathbb{N}_0\pi_a\cap R^{\mathrm{re}}_a$. Definition 5(6) implies that for each $i\in N$ and $a\in A$ one has $\sigma_{i,a}(R^{\mathrm{re}}_a)=R^{\mathrm{re}}_{i\triangleright a}$. Note that this definition of "real roots" coincides with the standard definition in case of Kac-Moody algebras. In general however our set R^{re}_a contains also "isotropic roots" (see examples below). The reason for this difference is the presence of a larger symmetry and the non-existence of a bilinear form.

Convention 6. For products of $\sigma_{i,a}$ we will use a similar convention as for products of $s_{i,a}$, see Convention 2.

Example 7. Let $(R,N,A,\triangleright)\in\mathcal{R}$ be of rank one. Then Definition 5(1) implies that A has cardinality 1 or 2. Further, by Definition 5(2),(4) one has $\pi_a=\{\alpha_a\}$ and $R_a=\{\alpha_a,-\alpha_a\}$ for each $a\in A$, where $\alpha_a\in\mathbb{R}\setminus\{0\}$. The other axioms do not give additional restrictions, and hence one has precisely two different elements in $\mathcal{R}$ which have rank 1.

Example 8. Let $A=\{a\}$ and $N=\{1,\dots,n\}$ for some $n\in\mathbb{N}$. The set A admits the trivial action $\triangleright$ of $F_2(N)$ given by $i\triangleright a:=a$ for all $i\in N$, $a\in A$. Let $\mathfrak{g}$ be a Kac-Moody algebra of rank n and let (R,π) and (R^{re},π) be the corresponding set of roots and real roots, respectively. Let $s_{i,a}:=s_i$ denote the reflection with respect to $\alpha_i\in\pi$, and define $S=\{s_1,\dots,s_n\}$. Then both $((R,\pi,S),N,A,\triangleright)$ and $((R^{\mathrm{re}},\pi,S),N,A,\triangleright)$ are in $\mathcal{R}$.

Example 9. Let $\mathfrak{g}$ be a finite dimensional contragredient Lie superalgebra of rank n and $\widetilde{R}$ its root system. Define $R=\widetilde{R}\setminus 2\widetilde{R}$ and $N=\{1,\dots,n\}$. Let $\{\pi_a=\{\alpha_{1,a},\dots,\alpha_{n,a}\}\,|\,a\in A\}$ be the set of all ordered bases of R, where A denotes an index set (that is each basis appears $n!$ times, each time with a different ordering). For any $i\in N$ and $a\in A$ let $r_{i,a}$ be the reflection on $\alpha_{i,a}$ if $\alpha_{i,a}$ is an even root, and the odd reflection if $\alpha_{i,a}$ is an odd root. By [Ser96, Lemma 6.4] there exists a unique $b\in A$ such that $r_{i,a}(\pi_a)=\pi_b$, $\alpha_{i,a}=-\alpha_{i,b}$, and $\alpha_{j,a}\in\alpha_{j,b}+\mathbb{N}_0\alpha_{i,b}$ for all $j\neq i$. Define $i\triangleright a:=b$ and $\sigma_{i,a}:=\mathrm{id}$. Then $\triangleright$ is an action of $F_2(N)$ on A, and all axioms (2)–(7) in Definition 5 are fulfilled. Let ψ denote the map from A to the set of all bases of R defined by ignoring the order. Let A' be an orbit of $\triangleright$ in A. By [Ser96, Lemma 6.4] $\psi|_{A'}$ is surjective and and by Lemma 21(iii) it is injective. By definition of A' one gets $(R',N,A',\triangleright)\in\mathcal{R}$, where $R'=\{(R_a,\pi_a,S_a)\,|\,a\in A'\}$. Note that the cardinality of A can be reduced further by identifying $a,b\in A'$ if $(\alpha_{i,a},\alpha_{j,a})=(\alpha_{i,b},\alpha_{j,b})$ and $\deg(\alpha_{i,a})=\deg(\alpha_{i,b})$ for all $i,j\in\{1,\dots,n\}$ (for a somewhat different example see also Section 7). Then Serganovas odd reflections in [Ser96, (1), (2) of Section 6] are reinterpreted in our setting as base changes in R.

Example 10. In [Hec06b] and [Hec04] arithmetic root systems are defined. They fit into the setting of Definition 5 as sketched below. These arithmetic root systems are closely related to Nichols algebras, and under some additional assumptions to quantized enveloping algebras. Finite arithmetic root systems are fully classified, see [Hec06a] and the references therein.

Let $\mathbf{k}$ be a field of characteristic zero, $n \in \mathbb{N}$, $E = \{\alpha_1, \dots, \alpha_n\}$ an ordered basis of $\mathbb{Z}^n$, and $q_{ij} \in \mathbf{k}\setminus\{0\}$, where $q_{ii} \neq 1$, for all $i, j \in \{1, \dots, n\}$. Let $\chi : \mathbb{Z}^n \times \mathbb{Z}^n \to \mathbf{k}\setminus\{0\}$ be the unique bicharacter such that $\chi(\alpha_i, \alpha_j) = q_{ij}$ for all $i, j \in \{1, \dots, n\}$. Assume now that for any $i \in \{1, \dots, n\}$ and any $j \neq i$ relation $(q_{ii}^m - 1)(q_{ii}^m q_{ij} q_{ji} - 1) = 0$ holds for some $m \in \mathbb{N}_0$, and denote the smallest such number by m_{ij}. Let $i \triangleright E$ denote the ordered basis of $\mathbb{Z}^n$ consisting of $i \triangleright \alpha_i = -\alpha_i$ and $i \triangleright \alpha_j := \alpha_i + m_{ij}\alpha_j$ for $j \neq i$, in the ordering induced by the one of E. Proceed inductively with the new bases, using now the values of the bicharacter χ instead of the q_{ij}. The assumption that the bases $i_1 \triangleright (i_2 \triangleright \cdots (i_m \triangleright E))$ always exist is crucial to obtain the setting in Definition 5, and no general algebraic condition on the q_{ij} is known which is equivalent to this property. Let A denote the set of all ordered bases of $\mathbb{Z}^n$ of the form $i_1 \triangleright (i_2 \triangleright \cdots (i_m \triangleright E))$. By definition, $\triangleright$ is a transitive action of $F_2(\{1, \dots, n\})$ on A.

The Nichols algebra attached to the matrix (q_{ij}) is $\mathbb{Z}^n$-graded, where the generators have degree α_i, and $i \in \{1, \dots, n\}$. Moreover it has a restricted PBW basis [Kha99] consisting of homogeneous elements, and the degrees of these PBW generators can be regarded as the set of positive roots Δ^+ with respect to the basis E. In particular, let $C = (c_{ij})_{i,j=1,\dots,n}$ be a symmetrizable Cartan matrix, and $d_i \in \mathbb{N}$ for $1 \le i \le n$ such that $(d_i c_{ij})$ is symmetric. Let $q \in \mathbf{k}$ be not a root of 1, and set $q_{ij} := q^{d_i c_{ij}}$. Then the Nichols algebra attached to the structure constants q_{ij} is exactly $U_q(\mathfrak{n}_+)$, where $\mathfrak{n}_+$ is the upper triangular part of the Kac-Moody algebra associated to C, and the set Δ^+ is exactly the set of positive roots associated to C.

Let $\Delta := \Delta^+ \cup -\Delta^+$, $R_a := \Delta$ and $s_{i,a} :=$ id for all $a \in A$ (see above) and $i \in \{1, \dots, n\}$. Then $s_{i,a}$ satisfies the conditions in Definition 5(5). The remaining axioms follow either from the construction or from the theory of Nichols algebras, in particular from [Hec06b, Proposition 1].

In case of $U_q(\mathfrak{n}_+)$ it is sufficient to consider a one-element set A, but with our definition of A this is not the case. However, in general it is possible to reduce the number of elements of A by introducing the following equivalence relation. Call $a = \{\beta_1, \dots, \beta_n\}$ and $b = \{\gamma_1, \dots, \gamma_n\}$ equivalent, if $\chi(\beta_i, \beta_i) = \chi(\gamma_i, \gamma_i)$ and $\chi(\beta_i, \beta_j)\chi(\beta_j, \beta_i) = \chi(\gamma_j, \gamma_i)\chi(\gamma_i, \gamma_j)$ for all $i, j \in \{1, \dots, n\}$. Since the matrix of $s_{i,a}$ with respect to the bases a and $i \triangleright a$ depends only on the values of χ, it depends only on the equivalence class of a. A similar property holds for R_a, and hence one may replace A by the set of its equivalence classes. If q_{ii} is not a root of 1 for all i (in particular in the above setting related to $U_q(\mathfrak{n}_+)$) then with this identification one obtains exactly the setting in Example 8. A very concrete example of different type can be found in Section 7.

Remark 11. In view of the large amount of examples given above it seems very difficult to determine all elements of $\mathcal{R}$. On the other hand, Theorem 29 shows that the structure of the elements of $\mathcal{R}$ is not as complicated as it looks at first sight. We see the rich internal structure together with the difficulty of classification as the fascinating key features of our definition.

We continue with analyzing the structure of the elements of $\mathcal{R}$.

Lemma 12. *For any $a \in A$ and $i \in N$ one has* $\sigma_{i,a}(R_a^+ \setminus \{\alpha_{i,a}\}) = R_{i\triangleright a}^+ \setminus \{\alpha_{i,i\triangleright a}\}$.

Proof. This follows from Definition 5(3),(4),(5). □

Corollary 13. *Let $m \in \mathbb{N}$, $(i_1, \dots, i_m) \in N^m$, and $a \in A$. Let $b = i_1 \cdots i_m \triangleright a$ and $w := \sigma_{i_2} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}$. Then one has*

$$\sigma_{i_1,i_1\triangleright b} w(R^+_a) \cap -R^+_b =$$

$$\begin{cases} \sigma_{i_1,i_1\triangleright b}(w(R^+_a) \cap -R^+_{i_1\triangleright b}) \cup \{-\alpha_{i_1,b}\} & \textit{if } \alpha_{i_1,i_1\triangleright b} \in w(R^+_a) \\ \sigma_{i_1,i_1\triangleright b}((w(R^+_a) \cap -R^+_{i_1\triangleright b}) \setminus \{-\alpha_{i_1,i_1\triangleright b}\}) & \textit{otherwise, that is } -\alpha_{i_1,i_1\triangleright b} \in w(R^+_a). \end{cases}$$

Corollary 14. *Let $m \in \mathbb{N}_0$, $(i_1, \dots, i_m) \in N^m$, and $a \in A$. For all $r \le m$ define $a_r := i_r \cdots i_m \triangleright a$, and let $w := \sigma_{i_1} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}$. Then one has $|w(R^+_a) \cap -R^+_{a_1}| \in m - 2\mathbb{N}_0$. Moreover if $|w(R^+_a) \cap -R^+_{a_1}| = m$ then*

$$w(R^+_a) \cap -R^+_{a_1} = \{-\sigma_{i_1} \cdots \sigma_{i_{r-2}} \sigma_{i_{r-1},a_r}(\alpha_{i_r,a_r}) \,|\, 1 \le r \le m\}. \tag{10}$$

Let N and A be nonempty sets and $\triangleright$ an action of $F_2(N)$ on A. Let N' be a nonempty subset of N. Then there is a unique decomposition $A = \cup_p A_p$ of A into the disjoint union of nonempty subsets such that $F_2(N')$ acts transitively on A_p for all p. Let $\triangleright_p$ denote the restriction of $\triangleright$ to $F_2(N') \times A_p$.

The following lemma can be obtained immediately from Definition 5.

Lemma 15. *Let $(R = \{(R_a, \pi_a, S_a) \,|\, a \in A\}, N, A, \triangleright) \in \mathcal{R}$ and N' a nonempty subset of N. Let $A = \cup_p A_p$ be the above decomposition with respect to the action of $F_2(N')$. For any $a \in A$ set*

$$\pi'_a := \{\alpha_{n,a} \,|\, n \in N'\}, \qquad S'_a := \{\sigma_{n,a} \,|\, n \in N'\}, \qquad R'_{p,a} := \mathbb{Z}\pi'_a \cap R_a. \tag{11}$$

Then $(R'_p, N', A_p, \triangleright_p) \in \mathcal{R}$, where $R'_p = \{(R'_{p,a}, \pi'_a, S'_a) \,|\, a \in A_p\}$.

5. The rank two case

In this section let $(R, N = \{i, j\}, A, \triangleright) \in \mathcal{R}$ have rank two. We will give a description of R^+_a, where $a \in A$, in terms of elements of some π_b, $b \in A$. Set $d := |R^+_a| \in \mathbb{N} \cup \{\infty\}$.

Lemma 16. *Let $a \in A$, $m \in \mathbb{N}_0$, $(i_1, \dots, i_m) \in N^m$, and $w = \sigma_{i_1} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}$. If $m < d$ and $w(\alpha_{i,a}) \in -R^+_b$, where $b = i_1 \cdots i_m \triangleright a$, then $w(\alpha_{j,a}) \in R^+_b$.*

Proof. Assume to the contrary that $w(\alpha_{j,a}) \in -R^+_b$. Then $w(R^+_a) \subset -R^+_b$ which is a contradiction to Corollary 14 and the relation $m < d$. □

Lemma 17. *Let $a \in A$. For all $n \in \mathbb{N}_0$ set $i_{2n} = j$, $i_{2n+1} = i$, $j_{2n} = i$, $j_{2n+1} = j$, $a_n = i_n \cdots i_1 \triangleright a$, and $b_n = j_n \cdots j_1 \triangleright a$.*

(i) If $d \in \mathbb{N}$ then

$$R^+_a = \{\sigma_{i_1} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a_m}(\alpha_{i_{m+1},a_m}) \,|\, 0 \le m < d\}. \tag{12}$$

In this case one has $\alpha_{j,a} = \sigma_{i_1} \cdots \sigma_{i_{d-2}} \sigma_{i_{d-1},a_{d-1}}(\alpha_{i_d,a_{d-1}})$.

(ii) If R_a is infinite then

$$\begin{aligned} (R^{\mathrm{re}}_a)^+ = \{\sigma_{i_1} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a_m}(\alpha_{i_{m+1},a_m}) \,|\, m \in \mathbb{N}_0\} \cup \\ \{\sigma_{j_1} \cdots \sigma_{j_{m-1}} \sigma_{j_m,b_m}(\alpha_{j_{m+1},b_m}) \,|\, m \in \mathbb{N}_0\}, \end{aligned} \tag{13}$$

and the elements given in Equation (13) are pairwise different from each other.

Proof. Set $w_0 := \mathrm{id}$, $w_1 := \sigma_{i,a}$, $M_0 := \emptyset$, and $M_1 := \{\alpha_{i,a}\}$. We inductively define $w_t \in \mathrm{End}(\mathbb{R}^2)$ and $M_t \subset R^+_a$ for $t \geq 0$ by the formulas

$$w_{t+1} := \sigma_{i_{t+1},a_t} w_t, \qquad M_{t+1} := M_t \cup \{w_t^{-1}(\alpha_{i_{t+1},a_t})\}. \tag{14}$$

We will show that

(1) $M_t = \{\beta \in R^+_a \mid w_t(\beta) \in -R^+_{a_t}\}$,
(2) $|M_t| = t$.

Setting $t = d$ in the above formulas and using the inductive definition of M_d one obtains Equation (12). Similarly, Equation (13) follows from the definition of R^{re}_a and Definition 5(6). The remaining statements of (i) and (ii) follow from the fact that $\alpha_{j,a} \notin M_t$ for $t < d$, which itself is a consequence of the assumption $\alpha_{i,a} \in M_t$ and Lemma 16.

We perform the proof of (1) and (2) by induction on t. Note that for $t = 0$ and $t = 1$ these formulas hold trivially. Assume now that M_t satisfies (1) and (2), and that relation $t < d$ holds. Then $w_t^{-1}(\alpha_{i_t,a_t}) = w_{t-1}^{-1}\sigma_{i_t,a_{t-1}}^{-1}(\alpha_{i_t,a_t}) = -w_{t-1}^{-1}(\alpha_{i_t,a_{t-1}}) \in -M_{t-1}$, so $w_t^{-1}(\alpha_{i_t,a_t}) \in -R^+_a$. Hence $w_t^{-1}(\alpha_{i_{t+1},a_t}) \in R^+_a$ by Lemma 16. This gives $\alpha_{i_{t+1},a_t} \in w_t(R^+_a)$. By Corollary 13 one has

$$\begin{aligned} R^+_a \cap -w_{t+1}^{-1}(R^+_{a_{t+1}}) &= w_{t+1}^{-1}(w_{t+1}(R^+_a) \cap -R^+_{a_{t+1}}) \\ &= w_{t+1}^{-1}(\sigma_{i_{t+1},a_t}(w_t(R^+_a) \cap -R^+_{a_t}) \cup \{-\alpha_{i_{t+1},a_{t+1}}\}) \\ &= w_t^{-1}(w_t(R^+_a) \cap -R^+_{a_t}) \cup \{-w_{t+1}^{-1}(\alpha_{i_{t+1},a_{t+1}})\} \\ &= M_t \cup \{w_t^{-1}(\alpha_{i_{t+1},a_t})\} = M_{t+1}. \end{aligned} \tag{15}$$

Hence (1) holds. Moreover, by the second equality of (15) and Corollary 13 one can see that (2) is valid. □

Lemma 18. *If d is finite then for all $a \in A$ the relations*

$$\begin{aligned} &\sigma_i\sigma_j\sigma_i\cdots\sigma_j\sigma_{i,a} = \sigma_j\sigma_i\sigma_j\cdots\sigma_i\sigma_{j,a} \ (d \text{ factors}) \quad \text{if } d \text{ is odd}, \\ &\sigma_j\sigma_i\sigma_j\cdots\sigma_j\sigma_{i,a} = \sigma_i\sigma_j\sigma_i\cdots\sigma_i\sigma_{j,a} \ (d \text{ factors}) \quad \text{if } d \text{ is even} \end{aligned} \tag{16}$$

hold.

Proof. By Definition 5(6),(7) the assertion of the lemma is equivalent to the equation $(\sigma_i\sigma_j)^{d-1}\sigma_i\sigma_{j,a} = \mathrm{id}$. If d is even then applying Lemma 17(i) twice gives that

$$\begin{aligned} &(\sigma_i\sigma_j)^{d-1}\sigma_i\sigma_{j,a}(\alpha_{i,a}) = (\sigma_i\sigma_j)^{d/2}\sigma_{i,i\triangleright a}(\alpha_{i,i\triangleright a}) \\ &\qquad = (\sigma_i\sigma_j)^{d/2-1}\sigma_i\sigma_{j,a}(-\alpha_{i,a}) = \sigma_{i,i\triangleright a}(-\alpha_{i,i\triangleright a}) = \alpha_{i,a}, \\ &(\sigma_i\sigma_j)^{d-1}\sigma_i\sigma_{j,a}(\alpha_{j,a}) = -(\sigma_i\sigma_j)^{d-1}\sigma_{i,j\triangleright a}(\alpha_{j,j\triangleright a}) \\ &\qquad = -(\sigma_i\sigma_j)^{d/2-1}\sigma_i\sigma_{j,a}(\alpha_{j,a}) = (\sigma_i\sigma_j)^{d/2-1}\sigma_{i,j\triangleright a}(\alpha_{j,j\triangleright a}) = \alpha_{j,a}. \end{aligned}$$

For odd d the claim follows similarly. □

6. THE GENERAL CASE

In this section we use the results in the rank two case to show that to any $(R, N, A, \triangleright) \in \mathcal{R}$ one can attach a faithful representation of a Coxeter groupoid in a natural way.

Proposition 19. *Let* $(R, N, A, \triangleright) \in \mathcal{R}$*. For each* $i, j \in N$*, where* $i \neq j$*, and any* $a \in A$ *set* $m_{i,j;a} := |\mathbb{N}_0\{\alpha_{i,a}, \alpha_{j,a}\} \cap R_a|$*. Set* $V = \bigoplus_{a\in A} V_a$*, where* $V_a = V_0 = \mathbb{R}^{|N|}$ *(see Definition 5(2)), and let* $P_a : V \to V_a$ *and* $\iota_a : V_a \to V$ *be the canonical projection and canonical injection, respectively. Then the assignment* $\rho : e_a \mapsto \iota_a P_a$*,* $s_{i,a} \mapsto \iota_{i\triangleright a}\sigma_{i,a}P_a$*, gives a representation* (ρ, V) *of the Coxeter groupoid* $(W, N, A, \triangleright, \mathbf{m})$*. In particular for* W *one has* $e_a \neq 0$ *and* $s_{i,a} \neq 0$ *for all* $i \in N$ *and* $a \in A$*.*

Proof. By construction and by Definition 5(6) one obtains that Equations (3) are compatible with the definition of the representation. By Definition 5(7) it remains to show that if $m_{i,j;a}$ is finite then equation

$$\begin{aligned} &\sigma_i\sigma_j\sigma_i\cdots\sigma_j\sigma_{i,a} = \sigma_j\sigma_i\sigma_j\cdots\sigma_i\sigma_{j,a}\ (m_{i,j:a}\text{ factors}) \quad \text{if } m_{i,j;a} \text{ is odd,}\\ &\sigma_j\sigma_i\sigma_j\cdots\sigma_j\sigma_{i,a} = \sigma_i\sigma_j\sigma_i\cdots\sigma_i\sigma_{j,a}\ (m_{i,j;a}\text{ factors}) \quad \text{if } m_{i,j:a} \text{ is even} \end{aligned} \tag{17}$$

holds. The restrictions of the above equations to the space $\mathbb{R}\alpha_{i,a} \oplus \mathbb{R}\alpha_{j,a}$ are valid by Lemma 18 and Lemma 15. This means that for $x_{i,j;a} := (\sigma_i\sigma_j)^{m_{i,j;a}-1}\sigma_i\sigma_{j,a}$ one has

$$x_{i,j;a}(\alpha_{i,a}) = \alpha_{i,a}, \quad x_{i,j;a}(\alpha_{j,a}) = \alpha_{j,a}, \quad x_{i,j;a}(\alpha_{n,a}) \in \alpha_{n,a} + \mathbb{N}_0\{\alpha_{i,a}, \alpha_{j,a}\}$$

for all $n \in N \setminus \{i, j\}$. Since $x_{i,j;a}(R_a) = R_a$ by Definition 5(5),(7), this implies that $x_{i,j;a}(\alpha_{n,a}) = \alpha_{n,a}$ for all $n \in N$. □

If $(R, N, A, \triangleright) \in \mathcal{R}$ then let $(\rho(W), R, N, A, \triangleright)$ denote the groupoid obtained as the image of the representation ρ in Proposition 19.

In the following lemmata let $(R, N, A, \triangleright) \in \mathcal{R}$, and for all $a \in A$ and $i, j \in N$, where $i \neq j$, set $m_{i,j;a} := |\mathbb{N}_0\{\alpha_{i,a}, \alpha_{j,a}\} \cap R_a|$. Note that one has

$$m_{i,j;a} = m_{j,i;a} = m_{i,j;i\triangleright a} = m_{i,j;j\triangleright a}. \tag{18}$$

We will analyze the properties of the Coxeter groupoid $(W, N, A, \triangleright, \mathbf{m})$.

Lemma 20. *Let* $m \in \mathbb{N}_0$*,* $(i_1, \dots, i_m) \in N^m$*, and* $a \in A$*. If* $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}) < m$ *then* $s_{i_2}\cdots s_{i_{m-1}}s_{i_m,a} = s_{i_1}s_{j_2}\cdots s_{j_{m-2}}s_{j_{m-1},a}$ *for some* $j_2, \dots, j_{m-1} \in N$*.*

Proof. By the definition of ℓ, the relation $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}) < m$ implies that equation $s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a} = s_{j_2}\cdots s_{j_{r-1}}s_{j_r,a}$ holds for some $r \leq m$. Moreover W is $\mathbb{Z}/2\mathbb{Z}$-graded, and hence m is even if and only if r is odd. Using Equations (3) one can assume that $r = m - 1$. Then the claim follows from the last equation in (3). □

Lemma 21. *Let* $m \in \mathbb{N}$*,* $(i_1, \dots, i_m) \in N^m$*, and* $a \in A$*.*

(i) If $\sigma_{i_1}\cdots\sigma_{i_{m-2}}\sigma_{i_{m-1},a}(\alpha_{i_m,a}) \in -R^+_b$*, where* $b = i_1\cdots i_{m-1} \triangleright a$*, then relation* $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,i_m\triangleright a}) < m$ *holds.*

(ii) If $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}) = m$*, and there exist* $k \in N\setminus\{i_1\}$*,* $\alpha \in \pi_a$*, and* $h_1, h_2 \in \mathbb{N}$ *such that* $\sigma_{i_2}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha) = h_1\alpha_{i_1,f} + h_2\alpha_{k,f}$*, where* $f = i_2\cdots i_m \triangleright a$*, then equation* $s_{i_2}\cdots s_{i_{m-1}}s_{i_m,a} = s_k s_{j_2}\cdots s_{j_{m-2}}s_{j_{m-1},a}$ *holds for some* $j_2, \dots, j_{m-1} \in N$*.*

(iii) One has $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}) = |\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(R^+_a) \cap -R^+_{i_1\cdots i_m\triangleright a}|$*.*

Proof. We prove all statements parallelly by induction on m. Note that for $m = 1$ the claim holds by Definition 5. Assume now that the lemma holds for $0, 1, \dots, m-1$.

To (i): By Equation (6) one can assume that $\ell(s_{i_2}\cdots s_{i_{m-1}}s_{i_m,i_m\triangleright a}) = m - 1$. Moreover the last equation in (3) yields that it is sufficient to consider the case

$i_1 \neq i_2$. By the converse of (i) for $m-1$ one has $\sigma_{i_2} \cdots \sigma_{i_{m-2}} \sigma_{i_{m-1},a}(\alpha_{i_m,a}) \in R^+_{i_1 \triangleright b}$. By Lemma 12, since $\sigma_{i_1,i_1 \triangleright b}(\alpha_{i_1,i_1 \triangleright b}) = -\alpha_{i_1,b}$, one has

$$\sigma_{i_2} \cdots \sigma_{i_{m-2}} \sigma_{i_{m-1},a}(\alpha_{i_m,a}) = \alpha_{i_1,i_1 \triangleright b}. \tag{19}$$

We prove now by induction, with letting $d := m_{i_1,i_2;b} \in (2+\mathbb{N}_0) \cup \{\infty\}$, that

$$s_{i_1} s_{i_2} \cdots s_{i_{m-1}} s_{i_m,i_m \triangleright a} = s_{i_1} s_{i_2} s_{i_{3 (\mathrm{mod}\ 2)}} \cdots s_{i_{d (\mathrm{mod}\ 2)}} s_{j_{d+1}} \cdots s_{j_{m-1}} s_{i_m,i_m \triangleright a} \tag{20}$$

for some $j_{d+1} \cdots j_{m-1} \in N$, where $n (\mathrm{mod}\ 2) \in \{1,2\}$ for $n \leq d$. We also show that $m > d$, and that for these $j_{d+1}, \dots, j_{m-1}$ relation

$$\sigma_{j_{d+1}} \cdots \sigma_{j_{m-2}} \sigma_{j_{m-1},a}(\alpha_{i_m,a}) = \alpha_{i_{d+1 (\mathrm{mod}\ 2)},a'}, \tag{21}$$

holds, where $a' = j_{d+1} \cdots j_{m-1} \triangleright a$. Thus in our setting we may apply Equation (4) to get

$$s_{i_1} \cdots s_{i_{m-1}} s_{i_m,i_m \triangleright a} = s_{i_2} s_{i_1} s_{i_2} \cdots s_{i_{d+1 (\mathrm{mod}\ 2)}} s_{j_{d+1}} \cdots s_{j_{m-1}} s_{i_m,i_m \triangleright a}.$$

Since $\sigma_{i_{d+1 (\mathrm{mod}\ 2)}} \sigma_{j_{d+1}} \cdots \sigma_{j_{m-2}} \sigma_{j_{m-1},a}(\alpha_{i_m,a}) = \sigma_{i_{d+1 (\mathrm{mod}\ 2)},a'}(\alpha_{i_{d+1 (\mathrm{mod}\ 2)},a'})$ is contained in $-R^+_{i_{d+1 (\mathrm{mod}\ 2)} \triangleright a'}$ and because of $d \geq 2$ we may apply (i) to conclude that $\ell(s_{i_1} \cdots s_{i_{m-1}} s_{i_m,i_m \triangleright a}) < m$.

Note that equation $i_n = i_{n (\mathrm{mod}\ 2)}$ holds trivially for $n \leq 2$. Assume now that it also holds for $n \leq p$, where $2 \leq p < d$. Then Equation (19) and Lemma 17 imply that

$$\sigma_{i_{p+1}} \cdots \sigma_{i_{m-2}} \sigma_{i_{m-1},a}(\alpha_{i_m,a}) = \underbrace{\sigma_{i_{p (\mathrm{mod}\ 2)}} \sigma_{i_{p-1 (\mathrm{mod}\ 2)}} \cdots \sigma_{i_1} \sigma_{i_2,i_1 \triangleright b}}_{p-1 \text{ factors}}(\alpha_{i_1,i_1 \triangleright b})$$

is an element in $R^+_{i_{p+1} \cdots i_{m-1} \triangleright a} \cap \mathbb{N}_0 \{\alpha_{i_1,i_{p+1} \cdots i_{m-1} \triangleright a}, \alpha_{i_2,i_{p+1} \cdots i_{m-1} \triangleright a}\}$, but it is not in $\pi_{i_{p+1} \cdots i_{m-1} \triangleright a}$. In particular $m-1 \geq p+1$. Therefore induction hypothesis (ii) can be applied to the element $s_{i_p} s_{i_{p+1}} \cdots s_{i_{m-2}} s_{i_{m-1},a} \in W$ and taking $k := i_{p+1 (\mathrm{mod}\ 2)}$ and $\alpha := \alpha_{i_m,a}$. One obtains that $s_{i_{p+1}} \cdots s_{i_{m-2}} s_{i_{m-1},a} = s_{i_{p+1} (\mathrm{mod}\ 2)} s_{j'_{p+2}} \cdots s_{j'_{m-2}} s_{j'_{m-1},a}$ for some $j'_{p+2}, \dots, j'_{m-1} \in N$. Thus induction on p gives that Equation (20) holds and that $m-1 \geq d$.

Equation (21) follows from Equation (19) and Lemma 17 using (20).

To (iii): If $\ell(s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a}) < m$ then the claim follows from the induction hypothesis (iii) and Proposition 19. So one may assume that $\ell(s_{i_1} \cdots s_{i_{m-1}} s_{i_m,a}) = m$ and hence $\ell(s_{i_2} \cdots s_{i_{m-1}} s_{i_m,a}) = m-1$. Corollary 14 gives that

$$|\sigma_{i_1} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(R^+_a) \cap -R^+_{i_1 \cdots i_m \triangleright a}| \in m - 2\mathbb{N}_0.$$

Assume that (iii) does not hold. The induction hypothesis (iii) and Corollary 13 imply that $-\alpha_{i_1,i_1 \triangleright c} = \sigma_{i_2} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(\beta)$ for some $\beta \in R^+_a$, where $c = i_1 \cdots i_m \triangleright a$. Hence $\sigma_{i_m} \cdots \sigma_{i_3} \sigma_{i_2,i_1 \triangleright c}(\alpha_{i_1,i_1 \triangleright c}) \in -R^+_a$. Since (i) is already proven for m, we conclude that $\ell(s_{i_m} \cdots s_{i_2} s_{i_1,c}) < m$ which is a contradiction to the assumption and Equation (5).

To (ii): By (iii) and Corollary 14 one has $-\alpha_{i_1,i_1 \triangleright f} = \sigma_{i_1} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(\beta)$ for some $\beta \in R^+_a$, and hence $\sigma_{i_2} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(\beta) = \alpha_{i_1,f}$. Suppose that $\beta' := \sigma_{i_m} \cdots \sigma_{i_3} \sigma_{i_2,f}(\alpha_{k,f}) \in R^+_a$. Then $\sigma_{i_2} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(\beta') = \alpha_{k,f}$, and hence

$$\sigma_{i_2} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(h_1 \beta + h_2 \beta') = h_1 \alpha_{i_1,f} + h_2 \alpha_{k,f} = \sigma_{i_2} \cdots \sigma_{i_{m-1}} \sigma_{i_m,a}(\alpha).$$

Since $h_1 \beta + h_2 \beta' \notin \pi_a$ but $\alpha \in \pi_a$, this gives a contradiction. Therefore relation $\sigma_{i_m} \cdots \sigma_{i_3} \sigma_{i_2,f}(\alpha_{k,f}) \in -R^+_a$ holds, and hence $\ell(s_{i_m} \cdots s_{i_2} s_{k,k \triangleright f}) < m$ by (i). Equation (5) and the $\mathbb{Z}/2\mathbb{Z}$-grading of W give that $s_k s_{i_2} \cdots s_{i_{m-1}} s_{i_m,a} =$

$s_{j_2}\cdots s_{j_{m-2}}s_{j_{m-1},a}$ for some $j_2,\dots,j_{m-1}\in N$. Multiplying the last equation with s_k from the left gives the claim. $\square$

Corollary 22. *Let $m\in\mathbb{N}$, $(i_1,\dots,i_m,j)\in N^{m+1}$, and $a\in A$, and suppose that $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a})=m$. Then $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a})=m-1$ if and only if $\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a})\in -R^+_{i_1\cdots i_m\triangleright a}$. Equivalently, one has $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a})=m+1$ if and only if $\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a})\in R^+_{i_1\cdots i_m\triangleright a}$.*

Proof. Equation (5), Corollary 13, and Lemma 21(iii) give that $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a})\in\{m-1,m+1\}$. Therefore the if part follows from Lemma 21(i). On the other hand, if $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a})=m-1$ then $\ell(s_j s_{i_m}\cdots s_{i_2}s_{i_1,b})=m-1$, where $b=i_1\cdots i_m\triangleright a$. Using the assumption $\ell(s_{i_m}\cdots s_{i_2}s_{i_1,b})=m$, Corollary 13 and Lemma 21(iii) give that $\sigma_{i_m}\cdots\sigma_{i_2}\sigma_{i_1,b}(\beta)=-\alpha_{j,a}$ for some $\beta\in R^+_b$. This implies the only if part of the claim. $\square$

Lemma 23. *Let $m\in\mathbb{N}$, $(i_0,i_1,\dots,i_m,j)\in N^{m+2}$, and $a\in A$, and suppose that $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a})=m$. If $\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a})=\alpha_{i_0,i_1\cdots i_m\triangleright a}$ then $i_0\neq i_1$ and*

$$s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}=s_{i_{1(\mathrm{mod}\ 2)}}s_{i_{2(\mathrm{mod}\ 2)}}\cdots s_{i_{d-1(\mathrm{mod}\ 2)}}s_{j_d}\cdots s_{j_{m-1}}s_{j_m,a},\tag{22}$$

$$\sigma_{j_d}\cdots\sigma_{j_{m-1}}\sigma_{j_m,a}(\alpha_{j,a})=\alpha_{i_{d(\mathrm{mod}\ 2)},j_d\cdots j_m\triangleright a}\tag{23}$$

for some $j_d,\cdots,j_m\in N$, where $d=m_{i_1,i_2;i_1\cdots i_m\triangleright a}$ and $k(\mathrm{mod}\ 2)\in\{0,1\}$ for $k\in\{1,\dots,d\}$.

Proof. By Corollary 22 one has $\ell(s_{i_1}\cdots s_{i_m}s_{j,j\triangleright a})=m+1$. By Lemma 21(iii) and Corollary 14, especially Equation (10), one has $-\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a})\neq -\alpha_{i_1,i_1\cdots i_m\triangleright a}$, so $i_1\neq i_0$. Since $\sigma_{i_0}\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m}(\alpha_{j,a})\in -R^+_{i_0\cdots i_m\triangleright a}$, one can apply the proof of Lemma 21(i), especially Equations (19), (20), and (21). $\square$

Theorem 24. *Let $(R,N,A,\triangleright)\in\mathcal{R}$, and for all $a\in A$ and $i,j\in N$, where $i\neq j$, set $m_{i,j;a}:=|\mathbb{N}_0\{\alpha_{i,a},\alpha_{j,a}\}\cap R_a|$. Then the representation (ρ,V) of the Coxeter groupoid $(W,N,A,\triangleright,\mathbf{m})$ is faithful.*

Proof. Let $a\in A$, $m\in\mathbb{N}_0$, and $(i_1,\dots,i_m)\in N^m$. One has to show that if $\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}=\mathrm{id}$ and $i_1\cdots i_m\triangleright a=a$ then $s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}=e_a$. This follows from Lemma 21(iii). $\square$

Remark 25. Theorem 24 gives one possible answer to Serganova's question about the analog of Weyl group mentioned in the introduction. On the one hand we do not consider the group proposed by Serganova. On the other hand Theorem 24 shows that with our definition one obtains a Coxeter groupoid attached to any contragredient Lie superalgebra, which acts faithfully on the root system, and contains all features of ordinary and odd reflections.

Corollary 26. *Let $(R,N,A,\triangleright)\in\mathcal{R}$ and assume that $d:=|R_a|$ is finite for one (that is each) $a\in A$. Then A is finite. Further, let $(W,N,A,\triangleright,\mathbf{m})$ be the associated Coxeter groupoid. Then for each $a\in A$ there exists a unique $w_a\in W$ such that $w_ae_a=w_a$ and $\ell(w_a)$ is maximal. Moreover, one has $\ell(w_a)=d$.*

Proof. If R_a is finite then $|R_a|$ is independent of $a\in A$ by Definition 5(5).

Corollary 22 implies that if $\ell(w_a)$ is maximal then $w_a(\alpha_{j,a})\in -R^+_b$ for all $j\in N$, where $b\in A$ is such that $e_bw_a=w_a$. Assume now that w_a and w'_a have maximal length. Then $w_a(w'_a)^{-1}$ maps R^+_c to R^+_b, where $b,c\in A$ are such that $e_bw_a=w_a$ and $e_cw'_a=w'_a$. By Lemma 21(iii) one gets $\ell(w_a(w'_a)^{-1})=0$, so $b=c$ and

$w_a(w'_a)^{-1}=e_b$. Hence $w_a=w'_a$. The existence of a w_a with maximal length follows also from Lemma 21(iii). Now the finiteness of A follows from the transitivity of $\triangleright$ and the fact that there are only finitely many elements $w\in W$ with $we_a=w$ and $\ell(w)\le |R^+_a|$. $\square$

Theorem 27 (weak exchange condition)**.** *Let* $m\in\mathbb{N}$, $(i_0,i_1,\dots,i_m,j)\in N^{m+2}$, *and* $a\in A$, *and suppose that* $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a})=m$. *If* $\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j,a})=\alpha_{i_0,i_1\cdots i_m\triangleright a}$ *then there exist* $r\in\mathbb{N}$, $(j_1,\dots,j_r)\in N^r$, $(k_1,\dots,k_{r+1})\in N^{r+1}$, *and* $(a_1,\dots,a_r)\in A^r$, *such that the following relations hold.*

$$s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}=C_{j_1,k_1;a_1}\cdots C_{j_r,k_r;a_r},\tag{24}$$

$$C_{j_1,k_1;a_1}\cdots C_{j_r,k_r;a_r}s_{j,j\triangleright a}=s_{i_0}C_{j_1,k_1;k_2\triangleright a_1}\cdots C_{j_r,k_r;k_{r+1}\triangleright a_r},\tag{25}$$

$$C_{j_1,k_1;k_2\triangleright a_1}\cdots C_{j_r,k_r;k_{r+1}\triangleright a_r}s_{j,a}=s_{i_0}C_{j_1,k_1;a_1}\cdots C_{j_r,k_r;a_r},\tag{26}$$

$$j_1=i_1,\ k_1=i_0,\ k_{r+1}=j,\ a_r=a,\ \sum_{t=1}^{r}m_{j_t,k_t;a_t}-r=m,\tag{27}$$

$$j_n\ne k_n \textit{ for } 1\le n\le r,\tag{28}$$

$$k_{n+1}=\begin{cases} j_n & \textit{if } m_{j_n,k_n;a_n} \textit{ is odd,}\\ k_n & \textit{if } m_{j_n,k_n;a_n} \textit{ is even,}\end{cases}\quad \textit{for } 1\le n\le r,\tag{29}$$

and for $2\le n\le r$ *one has*

$$a_{n-1}=\begin{cases} j_nk_n\cdots j_nk_n\triangleright a_n\ \ (m_{j_n,k_n;a_n}-1 \textit{ factors}) & \textit{if } m_{j_n,k_n;a_n} \textit{ is odd,}\\ j_nk_n\cdots k_nj_n\triangleright a_n\ \ (m_{j_n,k_n;a_n}-1 \textit{ factors}) & \textit{if } m_{j_n,k_n;a_n} \textit{ is even.}\end{cases}\tag{30}$$

Proof. We show Equations (24),(25), and (26) in a way that (27), (28), (29), and (30) also hold.

Equation (24) is obtained easily by iterated application of Lemma 23. By Equations (8) and (27)-(30), for $1\le n\le r$, one has $C_{j_n,k_n;a_n}s_{k_{n+1},k_{n+1}\triangleright a_n}=s_{k_n}C_{j_n,k_n;k_{n+1}\triangleright a_n}$. Hence Equation (25) holds. Equation (26) can be obtained from (25) by multiplication with $s_{j,a}$ from the right and s_{i_0} from the left. $\square$

Now we are going to prove a generalization of Matsumoto's theorem [Mat64]. First we need a definition.

Definition 28. Let $(W,N,A,\triangleright,\mathbf{m})$ be a Coxeter groupoid. Let $\widetilde{W}$ denote the semigroup generated by the set $\{0,\tilde e_a,\tilde s_{i,a}\,|\,a\in A, i\in N\}$ and the relations

$$00=0,\quad 0\tilde e_a=\tilde e_a0=0\tilde s_{i,a}=\tilde s_{i,a}0=0,\tag{31}$$

$$\tilde e_a^2=\tilde e_a,\quad \tilde e_a\tilde e_b=0 \text{ for } a\ne b,\quad \tilde e_{i\triangleright a}\tilde s_{i,a}=\tilde s_{i,a}\tilde e_a=\tilde s_{i,a},\tag{32}$$

$$\begin{array}{ll}\tilde s_i\tilde s_j\cdots\tilde s_j\tilde s_{i,a}=\tilde s_j\tilde s_i\cdots\tilde s_i\tilde s_{j,a}\ (m_{i,j;a}\text{ factors}) & \text{if } m_{i,j;a} \text{ is finite and odd,}\\ \tilde s_j\tilde s_i\cdots\tilde s_j\tilde s_{i,a}=\tilde s_i\tilde s_j\cdots\tilde s_i\tilde s_{j,a}\ (m_{i,j;a}\text{ factors}) & \text{if } m_{i,j;a} \text{ is finite and even,}\end{array}\tag{33}$$

where a similar convention as in Convention 2 is used. More precisely we will write $(\widetilde{W},N,A,\triangleright,\mathbf{m})$ for this semigroup.

Analogously to Equation (7) we define

$$\tilde C_{i,j;a}=\begin{cases}\tilde s_i\tilde s_j\tilde s_i\cdots\tilde s_i\tilde s_{j,a}\ (m_{i,j;a}-1\text{ factors}) & \text{if } m_{i,j;a} \text{ is odd,}\\ \tilde s_i\tilde s_j\tilde s_i\cdots\tilde s_j\tilde s_{i,a}\ (m_{i,j;a}-1\text{ factors}) & \text{if } m_{i,j;a} \text{ is even.}\end{cases}\tag{34}$$

Similarly to Equations (8),(25) and (26) we have

$$\tilde{s}_i\tilde{C}_{j,i;a} = \begin{cases} \tilde{C}_{j,i:j\triangleright a}\tilde{s}_{j,a} & \text{if } m_{j,i;a} \text{ is odd,} \\ \tilde{C}_{j,i:i\triangleright a}\tilde{s}_{i,a} & \text{if } m_{j,i;a} \text{ is even,} \end{cases} \tag{35}$$

$$\tilde{C}_{j_1,k_1;a_1}\cdots\tilde{C}_{j_r,k_r;a_r}\tilde{s}_{j,j\triangleright a} = \tilde{s}_{i_0}\tilde{C}_{j_1,k_1;k_2\triangleright a_1}\cdots\tilde{C}_{j_r,k_r;k_{r+1}\triangleright a_r}, \tag{36}$$

$$\tilde{C}_{j_1,k_1;k_2\triangleright a_1}\cdots\tilde{C}_{j_r,k_r;k_{r+1}\triangleright a_r}\tilde{s}_{j,a} = \tilde{s}_{i_0}\tilde{C}_{j_1,k_1;a_1}\cdots\tilde{C}_{j_r,k_r;a_r}, \tag{37}$$

where i, j, j_n, k_n, a_n are as in Theorem 27.

Theorem 29. *Let* $(R, N, A, \triangleright) \in \mathcal{R}$, *and for all* $a \in A$ *and* $i, j \in N$, *where* $i \neq j$, *set* $m_{i,j;a} := |\mathbb{N}_0\{\alpha_{i,a}, \alpha_{j,a}\} \cap R_a|$. *Suppose that* $m \in \mathbb{N}_0$, $a \in A$, *and* $(i_1, \ldots, i_m), (j_1, \ldots, j_m) \in N^m$ *such that* $\ell(s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}) = m$ *and equation* $s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a} = s_{j_1}\cdots s_{j_{m-1}}s_{j_m,a}$ *holds in* $(W, N, A, \triangleright, \mathbf{m})$. *Then in the semigroup* $(\widetilde{W}, N, A, \triangleright, \mathbf{m})$ *one has* $\tilde{s}_{i_1}\cdots\tilde{s}_{i_{m-1}}\tilde{s}_{i_m,a} = \tilde{s}_{j_1}\cdots\tilde{s}_{j_{m-1}}\tilde{s}_{j_m,a}$.

Proof. We proceed by induction on m, where for $m = 0$ and $m = 1$ the claim holds by Proposition 19 and Lemma 12.

Assume now that the claim holds for some $m \in \mathbb{N}$. Since $s_{i_1}\cdots s_{i_m}s_{j_m,j_m\triangleright a} = s_{j_1}\cdots s_{j_{m-2}}s_{j_{m-1},j_m\triangleright a}$, we have $\ell(s_{i_1}\cdots s_{i_m}s_{j_m,j_m\triangleright a}) = m - 1$. By Corollary 22 this implies that $\sigma_{i_1}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j_m,a}) \in -R^+_{i_1\cdots i_m\triangleright a}$. Therefore there exists $t \in \{1, \ldots, m\}$ such that relations $\sigma_{i_{t+1}}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j_m,a}) \in R^+_{i_{t+1}\cdots i_m\triangleright a}$ and $\sigma_{i_t}\sigma_{i_{t+1}}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j_m,a}) \in -R^+_{i_t i_{t+1}\cdots i_m\triangleright a}$ hold. Thus by Lemma 12 one has $\sigma_{i_{t+1}}\cdots\sigma_{i_{m-1}}\sigma_{i_m,a}(\alpha_{j_m,a}) = \alpha_{i_t,i_{t+1}\cdots i_m\triangleright a}$. Now we apply Theorem 27 and obtain that there exists $r \in \mathbb{N}$ and $(h_1, \ldots, h_r) \in N^r$ such that

$$s_{i_{t+1}}\cdots s_{i_{m-1}}s_{i_m,a} = C_{h_1,k_1;a_1}\cdots C_{h_r,k_r;a_r} \tag{38}$$

where the same notation as in Theorem 27 is used. Note that $k_1 = i_t$, $h_1 = i_{t+1}$, $k_{r+1} = j_m$, and $a_r = a$. By Equation (25) we obtain that

$$s_{i_1}\cdots s_{i_{m-1}}s_{i_m,a}s_{j_m,j_m\triangleright a} = s_{i_1}\cdots s_{i_{t-1}}C_{h_1,k_1;k_2\triangleright a_1}\cdots C_{h_r,k_r;k_{r+1}\triangleright a_r}. \tag{39}$$

By induction hypothesis this yields

$$\tilde{s}_{j_1}\cdots\tilde{s}_{j_{m-2}}\tilde{s}_{j_{m-1},j_m\triangleright a} = \tilde{s}_{i_1}\cdots\tilde{s}_{i_{t-1}}\tilde{C}_{h_1,k_1;k_2\triangleright a_1}\cdots\tilde{C}_{h_r,k_r;k_{r+1}\triangleright a_r}. \tag{40}$$

Since $t \geq 1$, induction hypothesis and Equation (38) give that

$$\tilde{s}_{i_{t+1}}\cdots\tilde{s}_{i_{m-1}}\tilde{s}_{i_m,a} = \tilde{C}_{h_1,k_1;a_1}\cdots\tilde{C}_{h_r,k_r;a_r} \tag{41}$$

By Equation (37) this implies that

$$\tilde{s}_{i_t}\cdots\tilde{s}_{i_{m-1}}\tilde{s}_{i_m,a} = \tilde{C}_{h_1,k_1;k_2\triangleright a_1}\cdots\tilde{C}_{h_r,k_r;k_{r+1}\triangleright a_r}\tilde{s}_{j_m,a}. \tag{42}$$

Now multiply Equation (40) from the right by $\tilde{s}_{j_m,a}$ and Equation (42) from the left by $\tilde{s}_{i_1}\cdots\tilde{s}_{i_{t-1}}$ to obtain the claim. □

7. An example

We demonstrate the structural results in the previous sections on an example related to Nichols algebras, see [Hec05, Table 2,row 15].

Let $A=\{a,b,c,d,e\}$ and $N=\{1,2,3\}$. Further, let $\pi_x:=\{\alpha_{1,x},\alpha_{2,x},\alpha_{3,x}\}$ be a basis of $\mathbb{Z}^3\subset\mathbb{R}^3$ for each $x\in A$. Define

$$\begin{aligned}
R^+_a:=\{&\alpha_{1,a},\alpha_{2,a},\alpha_{3,a},\alpha_{1,a}+\alpha_{2,a},\alpha_{2,a}+\alpha_{3,a},2\alpha_{2,a}+\alpha_{3,a},\alpha_{1,a}+\alpha_{2,a}+\alpha_{3,a},\\
&\alpha_{1,a}+2\alpha_{2,a}+\alpha_{3,a},\alpha_{1,a}+2\alpha_{2,a}+2\alpha_{3,a},\alpha_{1,a}+3\alpha_{2,a}+2\alpha_{3,a}\},\\
R^+_b:=\{&\alpha_{1,b},\alpha_{2,b},\alpha_{3,b},\alpha_{1,b}+\alpha_{2,b},\alpha_{2,b}+\alpha_{3,b},\alpha_{1,b}+2\alpha_{2,b},2\alpha_{2,b}+\alpha_{3,b},\\
&\alpha_{1,b}+\alpha_{2,b}+\alpha_{3,b},\alpha_{1,b}+2\alpha_{2,b}+\alpha_{3,b},\alpha_{1,b}+3\alpha_{2,b}+\alpha_{3,b}\},\\
R^+_c:=\{&\alpha_{1,c},\alpha_{2,c},\alpha_{3,c},\alpha_{1,c}+\alpha_{2,c},\alpha_{2,c}+\alpha_{3,c},\alpha_{1,c}+\alpha_{2,c}+\alpha_{3,c},\alpha_{1,c}+2\alpha_{2,c}+\alpha_{3,c},\\
&2\alpha_{1,c}+2\alpha_{2,c}+\alpha_{3,c},\alpha_{1,c}+2\alpha_{2,c}+2\alpha_{3,c},2\alpha_{1,c}+3\alpha_{2,c}+2\alpha_{3,c}\},\\
R^+_d:=\{&\alpha_{1,d},\alpha_{2,d},\alpha_{3,d},\alpha_{1,d}+\alpha_{2,d},\alpha_{2,d}+\alpha_{3,d},\alpha_{1,d}+2\alpha_{2,d},\alpha_{1,d}+\alpha_{2,d}+\alpha_{3,d},\\
&\alpha_{1,d}+2\alpha_{2,d}+\alpha_{3,d},2\alpha_{1,d}+2\alpha_{2,d}+\alpha_{3,d},2\alpha_{1,d}+3\alpha_{2,d}+\alpha_{3,d}\},\\
R^+_e:=\{&\alpha_{1,e},\alpha_{2,e},\alpha_{3,e},\alpha_{1,e}+\alpha_{2,e},\alpha_{1,e}+\alpha_{3,e},\alpha_{2,e}+\alpha_{3,e},\alpha_{1,e}+\alpha_{2,e}+\alpha_{3,e},\\
&2\alpha_{1,e}+\alpha_{2,e}+\alpha_{3,e},\alpha_{1,e}+\alpha_{2,e}+2\alpha_{3,e},2\alpha_{1,e}+\alpha_{2,e}+2\alpha_{3,e}\}.
\end{aligned}$$

According to the following figure



define the maps $\sigma_{i,x}$ for all $i\in\{1,2,3\}$ and $x\in A$. Edges between Dynkin diagrams $x,y\in A$ labelled by the number i mean that $i\triangleright x=y$ (and hence $i\triangleright y=x$). If there is no edge with label i which connects a given Dynkin diagram x with another diagram, then one has $i\triangleright x=x$. The numbers $m\in\mathbb{N}_0$ in the elements $\sigma_{i,x}(\alpha_{j,x})=\alpha_{j,y}+m\alpha_{i,y}$ are defined in the conventional way. Thus one has

$$\begin{aligned}
\sigma_{1,a}(\alpha_{1,a})&=-\alpha_{1,c}, & \sigma_{1,a}(\alpha_{2,a})&=\alpha_{2,c}+\alpha_{1,c}, & \sigma_{1,a}(\alpha_{3,a})&=\alpha_{3,c},\\
\sigma_{2,a}(\alpha_{2,a})&=-\alpha_{2,a}, & \sigma_{2,a}(\alpha_{1,a})&=\alpha_{1,a}+\alpha_{2,a}, & \sigma_{2,a}(\alpha_{3,a})&=\alpha_{3,a}+2\alpha_{2,a},\\
\sigma_{3,a}(\alpha_{3,a})&=-\alpha_{3,b}, & \sigma_{3,a}(\alpha_{1,a})&=\alpha_{1,b}, & \sigma_{3,a}(\alpha_{2,a})&=\alpha_{2,b}+\alpha_{3,b},
\end{aligned}$$

$$\begin{aligned}
\sigma_{1,b}(\alpha_{1,b})&=-\alpha_{1,d}, & \sigma_{1,b}(\alpha_{2,b})&=\alpha_{2,d}+\alpha_{1,d}, & \sigma_{1,b}(\alpha_{3,b})&=\alpha_{3,d},\\
\sigma_{2,b}(\alpha_{2,b})&=-\alpha_{2,b}, & \sigma_{2,b}(\alpha_{1,b})&=\alpha_{1,b}+2\alpha_{2,b}, & \sigma_{2,b}(\alpha_{3,b})&=\alpha_{3,b}+2\alpha_{2,b},\\
\sigma_{3,b}(\alpha_{3,b})&=-\alpha_{3,a}, & \sigma_{3,b}(\alpha_{1,b})&=\alpha_{1,a}, & \sigma_{3,b}(\alpha_{2,b})&=\alpha_{2,a}+\alpha_{3,a},
\end{aligned}$$

$$\begin{aligned}
\sigma_{1,c}(\alpha_{1,c})&=-\alpha_{1,a}, & \sigma_{1,c}(\alpha_{2,c})&=\alpha_{2,a}+\alpha_{1,a}, & \sigma_{1,c}(\alpha_{3,c})&=\alpha_{3,a},\\
\sigma_{2,c}(\alpha_{2,c})&=-\alpha_{2,e}, & \sigma_{2,c}(\alpha_{1,c})&=\alpha_{1,e}+\alpha_{2,e}, & \sigma_{2,c}(\alpha_{3,c})&=\alpha_{3,e}+\alpha_{2,e},\\
\sigma_{3,c}(\alpha_{3,c})&=-\alpha_{3,d}, & \sigma_{3,c}(\alpha_{1,c})&=\alpha_{1,d}, & \sigma_{3,c}(\alpha_{2,c})&=\alpha_{2,d}+\alpha_{3,d},
\end{aligned}$$

$$\begin{aligned}
\sigma_{1,d}(\alpha_{1,d}) =& -\alpha_{1,b}, & \sigma_{1,d}(\alpha_{2,d}) =&\alpha_{2,b}+\alpha_{1,b}, & \sigma_{1,d}(\alpha_{3,d}) =&\alpha_{3,b},\\
\sigma_{2,d}(\alpha_{2,d}) =& -\alpha_{2,d}, & \sigma_{2,d}(\alpha_{1,d}) =&\alpha_{1,d}+2\alpha_{2,d}, & \sigma_{2,d}(\alpha_{3,d}) =&\alpha_{3,d}+\alpha_{2,d},\\
\sigma_{3,d}(\alpha_{3,d}) =& -\alpha_{3,c}, & \sigma_{3,d}(\alpha_{1,d}) =&\alpha_{1,c}, & \sigma_{3,d}(\alpha_{2,d}) =&\alpha_{2,c}+\alpha_{3,c},
\end{aligned}$$

$$\begin{aligned}
\sigma_{1,e}(\alpha_{1,e}) =& -\alpha_{1,e}, & \sigma_{1,e}(\alpha_{2,e}) =&\alpha_{2,e}+\alpha_{1,e}, & \sigma_{1,e}(\alpha_{3,e}) =&\alpha_{3,e}+\alpha_{1,e},\\
\sigma_{2,e}(\alpha_{2,e}) =& -\alpha_{2,c}, & \sigma_{2,e}(\alpha_{1,e}) =&\alpha_{1,c}+\alpha_{2,c}, & \sigma_{2,e}(\alpha_{3,e}) =&\alpha_{3,c}+\alpha_{2,c},\\
\sigma_{3,e}(\alpha_{3,e}) =& -\alpha_{3,e}, & \sigma_{3,e}(\alpha_{1,e}) =&\alpha_{1,e}+\alpha_{3,e}, & \sigma_{3,e}(\alpha_{2,e}) =&\alpha_{2,e}+\alpha_{3,e}.
\end{aligned}$$

One can check that $\sigma_i(R_x) = R_{i\triangleright x}$ for all $i \in \{1,2,3\}$ and $x \in \{a,b,c,d,e\}$. Proposition 19 tells that the following relations hold in $(W, N, A, \triangleright, \mathbf{m})$.

$$\begin{aligned}
s_1s_2s_{1,a} =&s_2s_1s_{2,a}, & s_1s_{3,a} =&s_3s_{1,a}, & s_2s_3s_2s_{3,a} =&s_3s_2s_3s_{2,a},\\
s_1s_2s_1s_{2,b} =&s_2s_1s_2s_{1,b}, & s_1s_{3,b} =&s_3s_{1,b}, & s_2s_3s_2s_{3,b} =&s_3s_2s_3s_{2,b},\\
s_1s_2s_{1,c} =&s_2s_1s_{2,c}, & s_1s_{3,c} =&s_3s_{1,c}, & s_2s_3s_{2,c} =&s_3s_2s_{3,c},\\
s_1s_2s_1s_{2,d} =&s_2s_1s_2s_{1,d}, & s_1s_{3,d} =&s_3s_{1,d}, & s_2s_3s_{2,d} =&s_3s_2s_{3,d},\\
s_1s_2s_{1,e} =&s_2s_1s_{2,e}, & s_1s_3s_{1,e} =&s_3s_1s_{3,e}, & s_2s_3s_{2,e} =&s_3s_2s_{3,e}.
\end{aligned}$$

Consider now the following longest word in W.

$$\begin{aligned}
&s_{1,b}s_{2,b}s_{1,d}s_{3,c}s_{2,e}s_{3,e}s_{1,e}s_{3,e}s_{2,c}s_{1,a} =\\
&s_{1,b}s_{2,b}s_{1,d}s_{2,d}s_{3,c}s_{2,e}s_{1,e}s_{3,e}s_{2,c}s_{1,a} =\\
&s_{2,d}s_{1,b}s_{2,b}s_{1,d}s_{3,c}s_{2,e}s_{1,e}s_{3,e}s_{2,c}s_{1,a} =\\
&s_{2,d}s_{1,b}s_{2,b}s_{3,a}s_{1,c}s_{2,e}s_{1,e}s_{3,e}s_{2,c}s_{1,a} =\\
&s_{2,d}s_{1,b}s_{2,b}s_{3,a}s_{2,a}s_{1,c}s_{2,e}s_{3,e}s_{2,c}s_{1,a} =\\
&s_{2,d}s_{1,b}s_{2,b}s_{3,a}s_{2,a}s_{1,c}s_{3,d}s_{2,d}s_{3,c}s_{1,a} =\\
&s_{2,d}s_{1,b}s_{2,b}s_{3,a}s_{2,a}s_{1,c}s_{3,d}s_{2,d}s_{1,b}s_{3,a}.
\end{aligned}$$

The second and the last line are an example for the weak exchange condition. However, one has the relation

$$s_{1,b}s_{2,b}s_{1,d}s_{3,c}s_{2,e}s_{3,e}s_{1,e}s_{3,e}s_{2,c}s_{1,a} \neq s_{2,c}s_{1,a}s_{3,b}s_{2,b}s_{3,a}s_{1,c}s_{3,d}s_{2,d}s_{1,b}s_{3,a}$$

for the trivial reason $1 \triangleright b = d \neq e = 2 \triangleright c$. This gives a counterexample to the standard form of the exchange condition.

References

[AS98] N. Andruskiewitsch and H.-J. Schneider, *Lifting of quantum linear spaces and pointed Hopf algebras of order p^3*, J. Algebra **209** (1998), 658–691.

[AS05] ______, *On the classification of finite-dimensional pointed Hopf algebras*, Preprint math.QA/0502157 (2005).

[Bor88] R.E. Borcherds, *Generalized Kac-Moody algebras*, J. Algebra **115** (1988), R501–512.

[CP61] A.H. Clifford and G.B. Preston, *The algebraic theory of semigroups*, Mathematical surveys, vol. 7, Amer. Math. Soc., Providence, Rhode Island, 1961.

[Dri87] V.G. Drinfel'd, *Quantum groups*, Proceedings ICM 1986, Amer. Math. Soc., 1987, pp. 798–820.

[Hec04] I. Heckenberger, *Rank 2 Nichols algebras with finite arithmetic root system*, Preprint math.QA/0412458 (2004).

[Hec05] ______, *Classification of arithmetic root systems of rank 3*, Preprint math.QA/0509145 (2005).

[Hec06a] ______, *Classification of arithmetic root systems*, Preprint math.QA/0605795 (2006).

[Hec06b] ______, *The Weyl groupoid of a Nichols algebra of diagonal type*, Invent. Math. **164** (2006), 175–188.

[Jim86] M. Jimbo, *A q-analog of* $\mathcal{U}(gl(N+1))$ *Hecke algebras, and the Yang-Baxter equation*, Lett. Math. Phys. **11** (1986), 247–252.

[Kac77] V.G. Kac, *Lie superalgebras*, Adv. Math. **26** (1977), 8–96.

[Kac90] ———, *Infinite dimensional lie algebras*, Cambridge Univ. Press, 1990.

[Kha99] V. Kharchenko, *A quantum analog of the Poincaré–Birkhoff–Witt theorem*, Algebra and Logic **38** (1999), no. 4, 259–276.

[LN05] O. Loos and E. Neher, *Reflection systems and partial root systems*, Preprint (2005).

[Mat64] H. Matsumoto, *Générateurs et relations des groupes de weyl généralisés*, C. R. Acad. Sci. Paris **258** (1964), 3419–3422.

[Ser96] V. Serganova, *On generalizations of root systems*, Commun. Algebra **24** (1996), 4281–4299.

[Yam99] H. Yamane, *On defining relations of affine Lie superalgebra and affine quantized universal enveloping superalgebras*, Publ. RIMS Kyoto Univ. **35** (1999), no. 3, 321–390.

István Heckenberger, Mathematisches Institut, Universität Leipzig, Augustusplatz 10-11, 04109 Leipzig, Germany.

E-mail address: `heckenberger@math.uni-leipzig.de`

Hiroyuki Yamane, Department of Pure and Applied Mathematics, Graduate School of Information Science and Technology, Osaka University, Toyonaka 560-0043, Japan

E-mail address: `yamane@ist.osaka-u.ac.jp`